

# EPSDT

## EARLY PERIODIC SCREENING, DIAGNOSIS, AND TREATMENT RHODE ISLAND MEDICAID PEDIATRIC ORAL HEALTH SCHEDULE

	Infancy							Early Childhood						Middle Childhood						Adolescence									
	Newborn	3-5 days	By 1 Mo	2 Mo	4 Mo	6 Mo	9 Mo	12 Mo	18 Mo	24 Mo	30 Mo	3 Yrs	4 Yrs	5 Yrs	6 Yrs	7 Yrs	8 Yrs	9 Yrs	10 Yrs	11 Yrs	12 Yrs	13 Yrs	14 Yrs	15 Yrs	16 Yrs	17 Yrs	18 Yrs	19 Yrs	20 Yrs
X To be performed O Perform when clinically necessary ↔ Perform within indicated time frame																													
Clinical oral examination <sup>1</sup>						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Assess oral growth and development <sup>2</sup>						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Caries-risk assessment <sup>3</sup>						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Radiographic assessment <sup>4</sup>						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Prophylaxis and topical fluoride <sup>4,5</sup>						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	0	0	0	0	
Fluoride supplementation <sup>5</sup>						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Anticipatory guidance/counseling <sup>6</sup>						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Oral hygiene counseling <sup>7</sup>						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Dietary counseling <sup>8</sup>						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Injury prevention counseling <sup>9</sup>						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Counseling for nonnutritive habits <sup>10</sup>						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Counseling for speech/language development <sup>11</sup>						↔	↔		X	X	X	X	X	X															
Substance abuse assessment <sup>11</sup>														X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Counseling for intraoral/perioral piercing														X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Counseling for HPV Vaccination <sup>11</sup>																				X	X	X	X	X	X	X	X	X	
Assessment and treatment of developing malocclusion						↔	↔		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Assessment for pit and fissure sealants <sup>12</sup>								X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	
Assessment and /or removal of third molars <sup>11</sup>																					X	X	X	X	X	X	X	X	
Transition to adult dental care																													

**NOTE:** RlTe Smiles is a Rhode Island dental program for children that's designed to improve access to dental care. Children who have Medicaid coverage who were born on or after May 1, 2000 are eligible. For more information on RlTe Smiles, go to <http://www.eohhs.ri.gov/Consumer/DentalServices/ChildrenYoungAdults.aspx>



## Footnotes:

- 1- First examination at the eruption of the first tooth and no later than 12 months. Repeat every 6 months or as indicated by child's risk status/susceptibility to disease. Includes assessment of pathology and injuries.
- 2- By clinical examination.
- 3- Must be repeated regularly and frequently to maximize effectiveness.
- 4- Timing, selection, and frequency determined by child's history, clinical findings, and susceptibility to oral disease.
- 5- Consider when systemic fluoride exposure is suboptimal. Up to at least 16 years.
- 6- Appropriate discussion and counseling should be an integral part of each visit for care.
- 7- Initially, responsibility of parent; as child develops, jointly with parent; then, when indicated, only child.
- 8- At every appointment; initially discuss appropriate feeding practices, then the role of refined carbohydrates and frequency of snacking in caries development and childhood obesity.
- 9- Initially play objects, pacifiers, car seats; then learning to walk; then with sports and routine playing, including the importance of mouthguards.
- 10- At first discuss the need for additional sucking: digits vs pacifiers; then the need to wean from the habit before malocclusion or skeletal dysplasia occurs. For school-aged children and adolescent patients, counsel regarding any existing habits such as fingernail biting, clenching, or bruxism.
- 11- Referral to a Pediatrician or specialist, if necessary.
- 12- For caries-susceptible primary molars, permanent molars, premolars, and anterior teeth with deep pits and fissures; placed as soon as possible after eruption.